



Gerhard RF Krueger MD PhD Adjunct Professor

Department of Pathology & Laboratory Medicine 6431 Fannin Street
Houston, Texas 77030 USA

Home Address: 3 Lahn Street 50859 Cologne, Germany e-Mail:
gerhard.krueger@uth.tmc.edu

August 21, 2017

To
Kirk Kimmerling DDS, President/CEO
Kimmerling Holdings Group LLC
3698 Largent Way, Suite 202, Marietta, GA 30064 USA

Report and Documentation

**A New Siloxane Embedded Benzalkonium Chloride - Based Skin
Protectant (fiteBac Germicidal Skin Softening Gel) for Use Against
Insect Bites and Related Dermal Infections.**

Una silicona nueva integrada con cloruro de benzalconio, para protección básica de la piel (fiteBac gel germicida suave para piel) para uso en picaduras de insectos e infecciones dérmicas relacionadas.

AUTHORS

Gerhard R.F.Krueger*, Dharam V. Ablashi**, Babak Baban*** and Clara Niedworok****

*Department of Pathology and Laboratory Medicine, UT-Houston Medical School, Houston, Texas 77030, USA

** HHV-6 Foundation, Santa Barbara, California, USA

***Departments of Oral Biology, Dental Surgery and Neurology, Medical College, University of Georgia, Augusta GA 30912

**** Dental School, University of Duesseldorf, Germany

Key Words: Germicidal Hand Gel, Insect Bites, Dermal Infections

Gel Germicida para Piel, Picaduras de Insectos, Infecciones Dérmicas

Contact Address: gerhard.krueger@uth.tmc.edu

Comments: Local institutional funding, no conflict of interest of any author

ABSTRACT

FiteBac Skin Softening Gel, widely used as a cosmetic, contains the anti-infectious benzalkonium chloride embedded in siloxane quaternary ammonium compounds, but doesn't contain alcohol. It thus serves well for skin protection combined with local anti-infectious activity. As it does not penetrate the skin, no systemic toxic side effects are expected, and respectively, more than 3 years of practical use have not revealed any adverse reactions.

We have tested in a pilot study the usefulness of fiteBac Skin Softening Gel against skin lesions following insect bites of various nature as well as a few other ailments (e.g. fungus-related intertrigo). Of the 23 cases tested, all but one responded to the treatment with complete resolution of skin lesions within a few days up to a week. The only exception was a case of suspected tick bite with local arthritic symptoms. In the latter one, a combination of the fiteBac gel treatment with

subsequent local administration of ibuprofen gel also lead to complete remission of dermatitis and arthritis.

We thus recommend a wider use of this cosmetic/anti-infectious agent for prevention and treatment of insect bites to avoid subsequent infectious complications (such as, for instance, Lyme disease).

RESUMEN

El gel fiteBac suave para piel, ha sido ampliamente empleado como cosmético, contiene cloruro de benzalconio y está integrado con compuestos de amoniaco cuaternario en silicona como anti-infeccioso, pero no contiene alcohol. Por lo tanto, sirve para la protección de la piel combinado con una actividad local anti-infecciosa. Como ésta no penetra la piel, no se esperan efectos secundarios, y en más de tres años de su aplicación no se han revelado reacciones adversas.

Se ha probado en un estudio piloto la utilidad del gel suavizante de piel en lesiones causadas por piquetes de insectos de diversa naturaleza, así como en otros padecimientos. (ej. micosis-relacionada con intertrigo). De 23 casos, todos menos uno respondió al tratamiento con la resolución total de las lesiones en la piel en unos cuantos días a una semana. La única excepción fue un caso con sospecha de mordedura de garrapata y síntomas de artritis local. Este último, la combinación del tratamiento con gel fiteBac y aplicación posterior de gel de ibuprofeno ayudó a la completa remisión de dermatitis y artritis.

Por lo anterior, se recomienda un uso más amplio de este agente cosmético/ anti-infeccioso para la prevención y tratamiento de picaduras de insectos para evitar complicaciones infecciosas subsecuentes. (Como, por ejemplo, la enfermedad de Lyme).

INTRODUCTION

Insect bites and stings may lead to transmission of pathogenic organisms and, in severe cases, are even treated as a medical emergency (1-4). Many of these are accompanied by secondary exanthema of various kind (5). Treatment commonly is done by epinephrine, diphenhydramine and steroids. Oral antibiotics may be given for infected bite wounds (6,7). Next step is to obtain a definitive diagnosis and to initiate appropriate treatments, for instance for such diseases as Lyme disease, West Nile fever, malaria, tularemia, plague, typhus and many others. Considering the world-wide presence of stinging and biting insects and the respective frequency of associated diseases, such conventional medical intervention affords a major effort in man-power and financial expenses. There are 30,000 cases of Lyme disease alone reported to the CDC annually (cdc.gov). Preventive measures for

immediate treatment of such stings and bites, even by repellents to avoid such stinging insect would be a major advantage for general health care.

Common insect repellents recommended by CDC and EPA (cdc.gov; epa.gov) include DEET (N,N-diethyl-meta-toluamide), IR3535 ((3-[N-butyl-N-acetyl] aminopropionic acid, ethyl ester), Picaridin and a few others (6,7). Such substances, although effective as insect repellent, do not fight infections if a few bites have occurred despite of their use and thus transferred possible infections. It would be a major advantage, therefore, to have a substance which could combine both activities: insect repellent and anti infectious actions.

We are presenting preliminary data of a study using a newly designed skin protectant creme which combines all three activities (skin protection, repellent, anti-infectious), and thus help to fight secondary infective diseases caused by biting and stinging insect.

MATERIALS AND METHODS

We used **fiteBac Germicidal Hand Softening Gel (GHSG)**: fig. 1, kindly provided by KHG fiteBac Technology, 3698 Largent Way, Marietta, GA 30064). The contents of the gel contain two cosmetics: an elastomer (cross linked silicon polymers) as found in lipsticks, antiperspirants, deodorants, and hair products. The second compound is a cosmetic ester PPG (Crodamol) also found in personal care products which combines to the elastomer to form a large polymeric organosilicon compound. The list of ingredient chemicals is as follows: Benzalkonium Chloride, Cyclopentasiloxane, Dimethicone Crosspolymer, PPG-3 Benzyl Ether Ethylhexanoate, Octadecyldimethyl Trimethoxysilylpropyl Ammonium Chloride. The anti-infectious activity of quaternary ammonium salts (QAS) and siloxanes is well documented from its use as polyurethane coating (8-12). The fiteBac Germicidal Hand Softening gel (**GHSG**), in laboratory testing, showed activities against various bacteria, and suggestive activities against several viruses and fungi (e.g. candida) (13, 14 [Molinari]) besides its skin cosmetic protectant activities.

This pilot study for an initial screening of GHSG included 23 volunteers with various acute skin disorders (see table), ages 12-81 years, 12 of which were males and 11 females. All consented to participate after respective risk information (“informed consent”).

The gel was administered to the skin immediately after bites of various insects with early symptoms occurring (redening, edema, itching, bullous dermatitis). The course of local dermatitis was monitored by photography with follow-up for up to 3 years. Patients were interviewed for eventual late symptoms (fever, head ache, chronic exanthema, arthritis and others). Biting insects were identified - as far as possible - by inspection (see table) .

RESULTS

All of the 23 cases but one treated with GHS gel reacted positively. Of these, 10 cases lead to complete resolution within 2-3 days. These included acute dermal reactions caused by the common mosquito *Culex pipiens*; a somewhat delayed reaction in lesions with dermal edema and blisters occurred in bites from the Chinese tiger mosquito, *Aedes albopicta* also named *Stegomyia albopicta* (3 days

to 5 days). No additional bites were observed after the application of GHS gel for protection of uninvolved skin areas, and no side effects of the gel itself were recorded. One case with several mosquito bites and a 1 day delay in application of the gel afforded 5 days to completely resolve.

Excellent resolution within 2-3 days following GHS gel administration was found in four cases of intertrigo (intertriginous dermatitis, ID). ID develops in skin folds exposed to heat, moisture and friction (e.g. axilla, perineum, inframammary creases and others) and is commonly super-infected by candida organisms or bacteria. Conventional treatment is by drying agents, local antimicrobial/antifungal drugs and occasionally topical steroid steroids (personal hygiene notwithstanding and testing for diabetes mellitus).

It took generally longer for complete resolution in cases with tick bites which included more severe local lesions such as blisters, nodular infiltrates, ulceration and - in one case - general exanthema. In such cases, at least one week of local GHS gel administration appeared advised.

Other cases which afforded longer (at least one week) administration included such not clearly identified cases as neurodermitis, certain “warts” (suspicious for papilloma virus infection), and “dry and choppy hands” (in the words of the patient). At least one week of treatment, if necessary longer, were required.

In one case, GHS gel was inefficient to resolve the problem: the patient had a suspected tick bite at the finger (not having identified the tick) with subsequent edema, swelling of the finger, and arthritic pain. After a week’s treatment with GHS gel, only the additional application of a local anti rheumatic ointment (ibuprofen) resulted in complete resolution after a second week.

DISCUSSION

FiteBac Germicidal Hand Softening Gel (GHSG) is available as over-the-counter medication regulated and approved as non-prescription product defined in FDA’s Topical Antimicrobial Monograph (RR Stewart, Senior Regulatory Consultant, Technology Science Advisory Group, Washington DC 20036, March 4, 2016). It serves as general skin cosmetic and protectant combined with antimicrobial (bacteria, viruses, certain fungi) activities (13,14). GHSG has several advantages over other preparations commonly used for similar indications. 1) it is a non-alcohol based hand sanitizer thus avoiding negative side effects of alcohols (15-17). It also avoids possible toxic effects of one of its ingredients (benzalkonium chloride) by its embedding in silicon polymers. It reduces resorption of the toxic ingredient significantly (18) while protecting the skin and its local microenvironment.

Considering such advantages, we tested GHSG in a pilot study with 23 cases of dermatitis following following various insect bites and related conditions. In all but

one case, GHSG lead to complete remission of skin pathology caused by insect bites of various kind as well as in 4 cases of intertrigo (a candida- or bacteria-caused dermatitis). The only case in which GHSG was not sufficient was by a delayed application to a tick bite which already had caused arthritic complications. This case also resolved after addition of the anti rheumatic ibuprofen. Our data thus warrant a greater use of GHSG for treating insect bites and some common dermatitis cases in the elderly (19). No toxic clinical effects were observed clinically during a 3 year period when used as a general cosmetic, and laboratory testing has shown that the gel does not penetrate the skin such as to initiate systemic toxicity (18).

Our as well as other data referred to above warrant the tentative use of GHSG against skin affections by candida (e.g. in diabetics, and in babies such as diaper rash), by certain viruses (e.g. papilloma virus “warts” and herpes labialis/genitalis) and possibly to also alleviate the symptoms of shingles. Special targets for using the gel should include nursing homes for the elderly, boys scout- and military camps, as well as for travelers in subtropical and tropical countries.

Besides, GHS gel appears to be useful as insect repellent to avoid biting, and possibly also as general dermal antiseptic especially, as it is not an alcohol-based ointment and it avoids possible toxic side effects of resorbed benzalkonium chloride and common alcoholic preparations.

REFERENCES

1. Lopez M, Diaz G, Cruz A, Mejia S, Villareal C, Retamoza G, Lopez P, Moreno C, Salazar C. Adverse reactions to mosquito bites scholars from Monterrey, Nuevo Leon, Mexico. *Rev Allerg Mex* 2010; 57: 79-84
2. Vasievich MP, Villareal JD, Tomecki KJ. Got the travel bug? A review of common infections, infestations, bites, and stings among returning travelers. *Am J Clin Dermatol* 2016; 17: 451-462
3. Schlagenhauf P, Weld L, Goorhuis A, Gaunert P, Weber R, von Sonnenburg F, Lopez-Velez R, Jensenius M, Cramer JP, Field VK, Odolini S, Gkrania-Klotsas E, Chappius F, Malvy D, von Genderen PJ, Mockenhaupt F, Jaureguiberry F, Smith C, Beeching NJ, Ursing J, Rapp C, Parola P, Grobusch MP. Travel-associated infection in Europe (2008-12): an analysis of EuroTravNet longitudinal, surveillance data, and evaluation of the effect of the pre-travel consultation. *Lancet Infect Dis* 2015; 15: 55-64
4. Langley R, Mack K, Haileyesus T, Proescholdbell S, Annet JL. National estimates of non-canine bite and sting injuries treated in US hospital emergency departments, 2001-2010. *Wilderness Environ Med* 2014; 25: 14-23
5. Drago F, Ciccarese G, Gasparini G, Cogorno L, Javor S, Toniolo A, Broccolo F. Contemporary infectious exanthema: an update. *Future Microbial* 2016; online publ; for reprints reprints@futuremedicine.com

6. Kuri-Morales PA, Correa-Morales F, Gonzales-Acosta C, Sanchez-Tejeda G, Moreno-Garcia M, Davalos-Becerril E, Juarez-Franco MF, Benitez-Alva JL, Gonzalez-Roldan JF. Repellency of 29 synthetic and natural commercial topical insect repellents against *Aedes aegypti* (Diptera: Culicidae) in Central Mexico. *J Med Entomol* 2017; DOI 10.1093/jme/tjx076
7. Kitchen LW, Lawrence KL, Coleman RE. The role of the United States military in the development of vector control products, including insect repellents, insecticides, and bed nets. *J Vector Ecol* 2009; 34: 50-61
8. Pant RR, Rasley BT, Buckley JP, Lloyd CT, Cozzens RF, Santangelo PG, Wynne JH. Synthesis, mobility study and antimicrobial evaluation of novel self-spreading ionic silicone oligomers. *J Appl Polym Sci* 2007; 104: 2954-2964
9. Pant RR, Buckley JL, Fulmer PA, Wynne JH, McCluskey DM, Phillips JP. Hybrid siloxane epoxy coatings containing quaternary ammonium moieties. *J Appl Polym Sci* 2008; 110: 3080-3086
10. Harney MB, Pant RR, Fulmer PA, Wynne JH. Surface self-concentrating amphiphilic quaternary ammonium biocides as coating additives. *ACS Appl Materials Interfaces* 2009; 1: 39-41
11. Pant RR, Fulmer PA, Harney MB, Buckley JP, Wynne JH. Synthesis and biocidal efficacy of self-spreading polydimethylsiloxane oligomers possessing oxyethylene-functionalized quaternary ammoniums. *J Appl Polym Sci* 2009; 113: 2397-2403
12. Wynne JH, Fulmer PA, McCluskey DM, Mackey NM, Buchanan JP. Synthesis and development of a multifunctional self-decontaminating polyurethane coating. *ACS Appl Materials Interfaces* 2011; 3: 2005-2011
13. Baban B, Liu JY, Tay FR, Pashley DH. Use of a new, simple, laboratory method for screening the antimicrobial and antiviral properties of hand sanitizers. *Am J Dent* 2012; 25: 327-331
14. Molinari JA, Nelson P. FiteBac germicidal hand gel study. Research report *The Dental Advisor* 2012; 43: 1-2 (see also www.dentaladvisor.com)
15. Loeffler H, Kampf G. Hand disinfection: how irritant are alcohols? *J Hosp Infect* 2008; 70: 44-48
16. Hibbard JS. Analysis comparing the antimicrobial activity and safety of current antiseptic agents. A review. *J Infus Nurs* 2005; 28: 194-207
17. Gormley NJ, Bronstein AC, Rasimas JJ, Pao M, Wratney AT, Sun J, Austin HA, Suffredini AF. The rising incidence of intentional ingestion of ethanol-containing hand sanitizers. *Crit Care Med* 2012; 40: 290-294
18. Rangappa S, Ajjarapu S, Supriya B, Fatima T, Murthy SN. An in vitro maximum usage (MU_{ST}) study of a hand sanitizer containing benzalkonium chloride. 2017 submitted for publication.

19. Lichterfeld-Kottner A, Hahnel E, Blume-Peytavi U, Kuttner J. Systematic mapping review about costs and economic evaluations of skin conditions and diseases in the aged. *J Tissue Viability* 2017; 26: 3-19

CAPIONS FOR FIGURES

Figure 1: fiteBac Hand Sanitizer as Provided by KHG fiteBac Technology

Figure 2: *Aedes albopictus* dermatitis and treatment effects. a: mosquito and acute vesicular dermatitis. b,c: drying of vesicles and disappearance of lesion after treatment with GHGS.

Figure 3: *Culex pipiens* effects and treatment with GHS gel. Top: mosquito. a-c: acute and post-treatment lesion.

Figure 4: *Ixodes ricinus* tick bites and treatment. Top and a,b: tick and acute bite lesion with exanthema migraines. c: treatment effects with GHSG.

Figure 5: European biting spider (suspected *Wolf spider*) and treatment. Top and a: spider and acute lesion. b: residual slight skin pigmentation after treatment.

Table 1: Clinical Data Base for GHS (germicidal hand softening) Gel Treatment

Patient (Age/Sex) Result	Diagnosis	Cause	Treatment
Mosquito bites			
BK 79/F flare and edema for 2 days drying of bullae, complete resolution 1 week	itching flare, edema, epidermal bullae	Chinese tiger mosquito <i>Aedes albopictus</i> (fig. 2)	GHS for 5 days, one/day
DA 44/M complete resolution	erythema, itching	common mosquito <i>Culex sp.</i>	GHS for 2 days
MBK 81/F complete resolution	erythema, itching	common mosquito <i>Culex sp.</i> (Fig. 3)	GHS for 3 days one/day
GRK 80/M complete resolution	erythema, itching	common mosquito <i>Culex sp.</i>	GHS for 3 days one/day

KN 53/F complete resolution	edema, flare, itching	unidentified mosquito	GHS for 5 days one/day
PF 12/M complete resolution	edema, itching, dermal blisters	Chinese tiger mosquito <i>Aedes albopictus</i>	GHS for 3 days twice/day
LMN 23/F complete resolution	flare, itching	unidentified insect	GHS for 4 days one/day
KS 40/M protection	mosquito bite prevention	common mosquito <i>Culex sp.</i>	GHS to arms & legs one/day
DV 18/M resolution in 3 days	rash, edema, vesicles	multiple unidentified mosquitoes	GHS to arms & legs one/day
LN 22/F resolution	flare, nodule, itching arms & legs	common mosquito <i>Culex sp.</i>	GHS for 2 days locally on lesions

Ticks and other

GK 78/M flare resolved in 2 days nodule in 5 days	edema, erythema, nodule	<i>Ixodes ricinus</i> (fig. 4)	GHS for 3 days twice/day
CN 18/F resolution in 5 days	erythema, itching	unidentified tick	GHS for 6 days one/day
MBK 79/F resolution in 1 week	flare, itching, edema blisters	spider bite (fig. 5)	GHS for 5 days one/day

Other Skin Affections

RBK 57/M edema, not pain resolution resolution after 2nd week	edema, local arthritis (finger)	suspected tick bite	GHS for 4 days ibuprofen creme
GRFK 80/M resolution	intertrigo	suspected candida	GPS for 2 days
RK 56/M shrinking & disappearance papilloma	rapid growing skin papilloma	suspected papilloma virus	GPS for 1 week twice/day

KA 68/F resolution	onychomycosis finger nails, eczema	fungus (unidentified)	GHS for 2 weeks
GK 78/M resolution	intertrigo	suspected candida	GHS for 3 days
LN 21/F resolution after reddening	itchy rash, burning	suspected neurodermitis	GHS for 5 days
CSN 51/F smooth skin after initial flare	“dry and choppy hands”	unidentified	GHS for 1 week
GK 76/M wart drying in 4 days resolution in 1 week	“fast growing wart”	suspected papilloma virus	GHS for 1 week
BP 75/F resolution	intertrigo inframammary crease	unidentified	GHS for 3 days
RF 80/M resolution	intertrigo inguinal	Candida	GHS for 4 days then twice a week

Figure 1: fiteBac Hand Sanitizer as Provided by KHG fiteBac Technology



Figure 2. *Aedes albopictus* dermatitis and treatment effects



Figure 3: *Culex pipiens* effects and treatment with GHS gel



Figure 4: *Ixodes ricinus* tick bites and treatment



Figure 5: European biting spider (suspected *Wolf spider*) and treatment

